

On the Dimensions of Cyclic Symmetry Classes of Tensors

M. R. Darafsheh, M. R. Pournaki

Abstract

The dimensions of the symmetry classes of tensors, associated with a certain cyclic subgroup of \mathcal{S}_m which is generated by a product of disjoint cycles is explicitly given in terms of the generalized Ramanujan sum. These dimensions can also be expressed as the Euler φ -function and the Möbius function.

Keywords: Symmetry class of tensors, Ramanujan sum, Euler φ -function, Möbius function.

1995 Mathematics Subject Classification: Primary 20C30, 11L07; Secondary 15A69.

1 Introduction

Let V be an n -dimensional vector space over the complex field \mathbb{C} . Let $\otimes^m V$ be the m th tensor power of V and write $\nu_1 \otimes \cdots \otimes \nu_m$ for the decomposable tensor product of the indicated vectors. To each permutation σ in \mathcal{S}_m there corresponds a unique linear operator $P(\sigma) : \otimes^m V \rightarrow \otimes^m V$ determined by $P(\sigma)(\nu_1 \otimes \cdots \otimes \nu_m) = \nu_{\sigma^{-1}(1)} \otimes \cdots \otimes \nu_{\sigma^{-1}(m)}$. Let G be a subgroup of \mathcal{S}_m and let $I(G)$ be the set of all the irreducible complex characters of G . It follows from the orthogonality relations for characters that

$$\left\{ T(G, \chi) : \otimes^m V \rightarrow \otimes^m V \mid T(G, \chi) = \frac{\chi(1)}{|G|} \sum_{\sigma \in G} \chi(\sigma) P(\sigma); \chi \in I(G) \right\}$$

is a set of annihilating idempotents which sum to the identity. The image of $\otimes^m V$ under the $T(G, \chi)$ is called the *symmetry class of tensors* associated with G and χ and is denoted by $V_\chi^m(G)$. It is well known that

$$\dim V_\chi^m(G) = \frac{\chi(1)}{|G|} \sum_{\sigma \in G} \chi(\sigma) n^{c(\sigma)}, \quad (1)$$

where $c(\sigma)$ is the number of cycles, including cycles of length 1, in the disjoint cycle decomposition of σ (see [4]).

Several papers are devoted to calculating $\dim V_\chi^m(G)$ in a more closed form than (1). Cummings [2] in the case that G is a cyclic subgroup of \mathbb{S}_m generated by a cycle of length m gives a formula for $\dim V_\chi^m(G)$ in terms of the Euler φ -function and considers the case that G is isomorphic to a direct product of cyclic groups as well. In [3] when G is the dihedral group of order $2m$ is considered and a formula is given when G is equal to the whole group \mathbb{S}_m in [5] and [6]. In all cases $\dim V_\chi^m(G)$ involves certain functions of n .

In [7] there is a formula for calculating $\dim V_\chi^m(G)$ in the case that $G = \langle \pi_1 \rangle \cdots \langle \pi_p \rangle$, where π_i s, $1 \leq i \leq p$, are disjoint cycles in \mathbb{S}_m . This formula involves the Euler φ -function and Möbius function and is a modification of the formula given in [2]. In this case if the order of π_i is m_i , $1 \leq i \leq p$, then $G \cong \mathbb{Z}_{m_1} \times \cdots \times \mathbb{Z}_{m_p}$. It is mentioned in [2] and [7] that if two groups are isomorphic, then the dimensions of the symmetry classes of tensors associated with them need not be equal. For example if $G = \langle (12) \rangle$ and $H = \langle (12)(34) \rangle$ are considered as subgroups of \mathbb{S}_4 , then it is easy to calculate that $\dim V_{\chi_0}^4(G) = n^3(n+1)/2$ and $\dim V_{\chi_0}^4(H) = n^2(n^2+1)/2$ whereas $G \cong H \cong \mathbb{Z}_2$ and χ_0 is the identity character of \mathbb{Z}_2 .

Now it is a natural question to consider the cyclic group $G = \langle \pi_1 \dots \pi_p \rangle$ where the π_i s, $1 \leq i \leq p$, are disjoint cycles and ask about the dimension of $V_\chi^m(G)$, where $\chi \in \mathbf{I}(G)$. In this case if the order of π_i , $1 \leq i \leq p$, is m_i , then $G \cong \mathbb{Z}_{[m_1, \dots, m_p]}$, where $[m_1, \dots, m_p]$ denotes the least common multiple of the integers m_1, \dots, m_p . In this paper we obtain a formula for $\dim V_\chi^m(G)$ in the above case and this formula involves the generalized Ramanujan sum which itself involves the Euler φ -function and Möbius function. Therefore for the rest of this paper let $G < \mathbb{S}_m$ be of the form

$$G = \langle \pi_1 \dots \pi_p \rangle,$$

where π_i s, $1 \leq i \leq p$, are disjoint cycles in \mathbb{S}_m of certain orders, say, m_1, \dots, m_p , respectively. Since G is cyclic, therefore the irreducible characters of G are all linear and are of the form

$$\begin{aligned} \chi_h : G \rightarrow \mathbb{C}^*, \quad \chi_h((\pi_1 \dots \pi_p)^t) &= \exp\left(\frac{2\pi i h t}{[m_1, \dots, m_p]}\right), \\ 0 \leq t &\leq [m_1, \dots, m_p] - 1, \end{aligned}$$

so

$$\mathbf{I}(G) = \left\{ \chi_h : G \rightarrow \mathbb{C}^* \mid 0 \leq h \leq [m_1, \dots, m_p] - 1 \right\},$$

where $[m_1, \dots, m_p]$ denotes the least common multiple of the integers m_1, \dots, m_p . The symbol (m_1, \dots, m_p) denotes the greatest common divisor of m_1, \dots, m_p .

2 A Result About the Ramanujan Sum

The well known Ramanujan sum is

$$C_m(h) = \sum_{\substack{t=0 \\ (t, m)=1}}^{m-1} \exp\left(\frac{2\pi i h t}{m}\right),$$

where m is a positive integer and h is a nonnegative integer. Ramanujan proved that (see [1])

$$C_m(h) = \frac{\varphi(m)\mu(m/(m, h))}{\varphi(m/(m, h))},$$

where φ is the Euler φ -function, i.e., $\varphi(1) = 1$; for $m > 1$, $\varphi(m) =$ the number of positive integers less than m and relatively prime to m , and μ is the Möbius function, i.e., $\mu(1) = 1$, $\mu(m) = 0$ if $p^2|m$ for some prime number p , and $\mu(m) = (-1)^r$ if $m = p_1 \dots p_r$, where p_1, \dots, p_r are distinct prime numbers.

For our main result, we need to generalize the Ramanujan sum. It seems natural to us to generalize the Ramanujan sum as follows.

Definition 1 Let m_1, \dots, m_p be positive integers and let h be a nonnegative integer. Suppose $d_1|m_1, \dots, d_p|m_p$. The *generalized Ramanujan sum* denoted by $S(h; m_1, \dots, m_p; d_1, \dots, d_p)$ is defined by

$$S(h; m_1, \dots, m_p; d_1, \dots, d_p) = \sum_{\substack{t=0 \\ (t, m_1)=d_1 \\ \vdots \\ (t, m_p)=d_p}}^{[m_1, \dots, m_p]-1} \exp\left(\frac{2\pi i h t}{[m_1, \dots, m_p]}\right).$$

If the set $\{0 \leq t \leq [m_1, \dots, m_p] - 1 \mid (t, m_i) = d_i; 1 \leq i \leq p\}$ is empty, then we define $S(h; m_1, \dots, m_p; d_1, \dots, d_p) = 0$.

Remark 1 It is obvious that $S(h; m; 1) = C_m(h)$, and so the sum appearing in Definition 1 is a generalization of the Ramanujan sum.

In the following lemma we prove that the generalized Ramanujan sum defined in Definition 1 involves the Ramanujan sum.

Lemma 1 Let m_1, \dots, m_p be positive integers and let h be a nonnegative integer. Suppose $d_1 | m_1, \dots, d_p | m_p$ and set $m'_i = m_i/d_i$, $M_i = m_1 \dots m_p/m_i$, $M'_i = m'_1 \dots m'_p/m'_i$, $D_i = d_1 \dots d_p/d_i$ ($1 \leq i \leq p$) and

$$l = \frac{(M_1, \dots, M_p)}{(M'_1, \dots, M'_p)(D_1, \dots, D_p)}.$$

Then we have

$$S(h; m_1, \dots, m_p; d_1, \dots, d_p) = \begin{cases} \frac{1}{l} C_{[m'_1, \dots, m'_p]}(hl) & , \quad \text{if } \left(\frac{[d_1, \dots, d_p]}{d_i}, m'_i \right) = 1, \\ & \quad \quad \quad 1 \leq i \leq p \\ 0 & , \quad \text{otherwise.} \end{cases}$$

Proof. By Definition 1 and the fact that $\exp(2\pi i h t / [m_1, \dots, m_p])$ is a periodic function of t with period $[m_1, \dots, m_p]$ we have

$$\begin{aligned} S(h; m_1, \dots, m_p; d_1, \dots, d_p) &= \sum_{\substack{t=0 \\ (t, m_1) = d_1 \\ \vdots \\ (t, m_p) = d_p}}^{[m_1, \dots, m_p]-1} \exp\left(\frac{2\pi i h t}{[m_1, \dots, m_p]}\right) \\ &= \frac{1}{l} \sum_{\substack{t=0 \\ (t, m_1) = d_1 \\ \vdots \\ (t, m_p) = d_p}}^{l[m_1, \dots, m_p]-1} \exp\left(\frac{2\pi i h t}{[m_1, \dots, m_p]}\right). \end{aligned}$$

Now letting $t = [d_1, \dots, d_p]t'$, we obtain

$$\begin{aligned}
S(h; m_1, \dots, m_p; d_1, \dots, d_p) &= \frac{1}{l} \sum_{\substack{t'=0 \\ ([d_1, \dots, d_p]t', m_1) = d_1 \\ \vdots \\ ([d_1, \dots, d_p]t', m_p) = d_p}}^{[m'_1, \dots, m'_p]-1} \exp\left(\frac{2\pi i h l t'}{[m'_1, \dots, m'_p]}\right) \\
&= \frac{1}{l} \sum_{\substack{t'=0 \\ (([d_1, \dots, d_p]/d_1)t', m'_1) = 1 \\ \vdots \\ (([d_1, \dots, d_p]/d_p)t', m'_p) = 1}}^{[m'_1, \dots, m'_p]-1} \exp\left(\frac{2\pi i h l t'}{[m'_1, \dots, m'_p]}\right).
\end{aligned}$$

If $([d_1, \dots, d_p]/d_i, m'_i) = 1$ for all i , $1 \leq i \leq p$, then the set of all the t 's indexing the above summation is equal to the set of all the t 's such that $0 \leq t' \leq [m'_1, \dots, m'_p] - 1$ with conditions $(t', m'_i) = 1$, $1 \leq i \leq p$. And if there is an i for which $([d_1, \dots, d_p]/d_i, m'_i) \neq 1$ then the above sum is zero and therefore we obtain:

$$\begin{aligned}
S(h; m_1, \dots, m_p; d_1, \dots, d_p) &= \begin{cases} \frac{1}{l} \sum_{\substack{t'=0 \\ (t', m'_1) = 1 \\ \vdots \\ (t', m'_p) = 1}}^{[m'_1, \dots, m'_p]-1} \exp\left(\frac{2\pi i h l t'}{[m'_1, \dots, m'_p]}\right) & , \text{ if } \left(\frac{[d_1, \dots, d_p]}{d_i}, m'_i\right) = 1, \\ & 1 \leq i \leq p \\ 0 & , \text{ otherwise,} \end{cases} \\
&= \begin{cases} \frac{1}{l} \sum_{\substack{t'=0 \\ (t', [m'_1, \dots, m'_p]) = 1}}^{[m'_1, \dots, m'_p]-1} \exp\left(\frac{2\pi i h l t'}{[m'_1, \dots, m'_p]}\right) & , \text{ if } \left(\frac{[d_1, \dots, d_p]}{d_i}, m'_i\right) = 1, \\ & 1 \leq i \leq p \\ 0 & , \text{ otherwise,} \end{cases} \\
&= \begin{cases} \frac{1}{l} C_{[m'_1, \dots, m'_p]}(hl) & , \text{ if } \left(\frac{[d_1, \dots, d_p]}{d_i}, m'_i\right) = 1, \\ & 1 \leq i \leq p \\ 0 & , \text{ otherwise.} \quad \square \end{cases}
\end{aligned}$$

In some special cases the generalized Ramanujan sum is given in the following examples:

Example 1 $S(0; m; d) = C_{m/d}(0) = \frac{\varphi(m/d)\mu((m/d)/(m/d,0))}{\varphi((m/d)/(m/d,0))} = \varphi(m/d).$

Example 2 If $(h, m) = 1$; we have

$$S(h; m; d) = C_{m/d}(h) = \frac{\varphi(m/d)\mu((m/d)/(m/d, h))}{\varphi((m/d)/(m/d, h))} = \mu(m/d).$$

3 The Dimensions of Some Symmetry Classes of Tensors

In this section, as we mentioned earlier, the group $G = \langle \pi_1 \dots \pi_p \rangle$ is considered, where the π_i s, $1 \leq i \leq p$, are disjoint cycles in \mathcal{S}_m . Our aim is to calculate $\dim V_{\chi}^m(G)$, where $\chi \in \mathcal{I}(G)$, in terms of known functions. Our formula involves the generalized Ramanujan sum.

Theorem 1 *Let $G = \langle \pi_1 \dots \pi_p \rangle$, where the π_i s, $1 \leq i \leq p$, are disjoint cycles in \mathcal{S}_m of orders m_1, \dots, m_p , respectively, and let χ_h , $0 \leq h \leq [m_1, \dots, m_p] - 1$, be an irreducible complex character of G . Then*

$$\dim V_{\chi_h}^m(G) = \frac{n^{m-(m_1+\dots+m_p)}}{[m_1, \dots, m_p]} \sum_{\substack{d_1|m_1 \\ \vdots \\ d_p|m_p}} S(h; m_1, \dots, m_p; d_1, \dots, d_p) n^{d_1+\dots+d_p},$$

where $S(h; m_1, \dots, m_p; d_1, \dots, d_p)$ denotes the generalized Ramanujan sum.

Proof. According to (1) the dimension of $V_{\chi_h}^m(G)$ is

$$\frac{1}{[m_1, \dots, m_p]} \sum_{\sigma \in G} \chi_h(\sigma) n^{c(\sigma)}, \tag{2}$$

where $c(\sigma)$ denotes the number of cycles, including cycles of length 1, in the disjoint cycle decomposition of σ . But every $\sigma \in G$ is equal to $\sigma = (\pi_1 \dots \pi_p)^t$ for some t , $0 \leq t \leq [m_1, \dots, m_p] - 1$. Since π_1, \dots, π_p are disjoint, so are $(\pi_1 \dots \pi_p)^t = \pi_1^t \dots \pi_p^t$. Appealing to [7] we can obtain

$$c(\pi_1^t \dots \pi_p^t) = c(\pi_1^t) + \dots + c(\pi_p^t) + m - (m_1 + \dots + m_p).$$

Note that if $(t, m_i) = d$, then π_i^t has d cycles of length m_i/d and therefore $c(\pi_i^t) = d = (t, m_i)$. So we have

$$c(\pi_1^t \dots \pi_p^t) = (t, m_1) + \dots + (t, m_p) + m - (m_1 + \dots + m_p).$$

Hence according to (2) we have

$$\begin{aligned} \dim V_{\chi_h}^m(G) &= \frac{1}{[m_1, \dots, m_p]} \sum_{\sigma \in G} \chi_h(\sigma) n^{c(\sigma)} \\ &= \frac{1}{[m_1, \dots, m_p]} \sum_{t=0}^{[m_1, \dots, m_p]-1} \chi_h(\pi_1^t \dots \pi_p^t) n^{c(\pi_1^t \dots \pi_p^t)} \\ &= \frac{1}{[m_1, \dots, m_p]} \sum_{t=0}^{[m_1, \dots, m_p]-1} \exp\left(\frac{2\pi i h t}{[m_1, \dots, m_p]}\right) n^{(t, m_1) + \dots + (t, m_p) + m - (m_1 + \dots + m_p)} \\ &= \frac{n^{m - (m_1 + \dots + m_p)}}{[m_1, \dots, m_p]} \sum_{t=0}^{[m_1, \dots, m_p]-1} \exp\left(\frac{2\pi i h t}{[m_1, \dots, m_p]}\right) n^{(t, m_1) + \dots + (t, m_p)}. \end{aligned}$$

Now letting $(t, m_i) = d_i$, $1 \leq i \leq p$, we obtain

$$\begin{aligned} \dim V_{\chi_h}^m(G) &= \frac{n^{m - (m_1 + \dots + m_p)}}{[m_1, \dots, m_p]} \sum_{\substack{d_1 | m_1 \\ \vdots \\ d_p | m_p}} \left(\sum_{\substack{t=0 \\ (t, m_1) = d_1 \\ \vdots \\ (t, m_p) = d_p}}^{[m_1, \dots, m_p]-1} \exp\left(\frac{2\pi i h t}{[m_1, \dots, m_p]}\right) \right) n^{d_1 + \dots + d_p} \\ &= \frac{n^{m - (m_1 + \dots + m_p)}}{[m_1, \dots, m_p]} \sum_{\substack{d_1 | m_1 \\ \vdots \\ d_p | m_p}} S(h; m_1, \dots, m_p; d_1, \dots, d_p) n^{d_1 + \dots + d_p}. \quad \square \end{aligned}$$

Using Theorem 1 we obtain Theorems 1 and 2 of [2] in the following corollaries.

Corollary 1 *If G is a cyclic subgroup of \mathbb{S}_m generated by an m -cycle and χ is the identity character 1, then $\dim V_\chi^m(G) = (1/m) \sum_{d|m} \varphi(m/d)n^d$.*

Proof. Since $\chi = \chi_0$, by Example 1 and using Theorem 1 we obtain

$$\dim V_\chi^m(G) = \frac{n^{m-m}}{m} \sum_{d|m} S(0; m; d)n^d = \frac{1}{m} \sum_{d|m} \varphi(m/d)n^d. \quad \square$$

Remark 2 In Corollary 1, if $\dim V = n = 1$, then $\dim \otimes^m V = 1$, and so $\dim V_\chi^m(G) = 0$ or 1. So $(1/m) \sum_{d|m} \varphi(m/d) = 0$ or 1. But $(1/m) \sum_{d|m} \varphi(m/d) = 0$ is impossible, therefore $(1/m) \sum_{d|m} \varphi(m/d) = 1$ or $\sum_{d|m} \varphi(d) = m$, which is well known identity in number theory.

Corollary 2 *If G is a cyclic subgroup of \mathbb{S}_m generated by an m -cycle and χ is a primitive linear character, then $\dim V_\chi^m(G) = (1/m) \sum_{d|m} \mu(m/d)n^d$.*

Proof. We know that a linear character of a cyclic subgroup of \mathbb{S}_m is primitive if its value on a generator of the subgroup is a primitive m th root of unity, so $\chi = \chi_h$ where $(h, m) = 1$ and by Example 2 and using Theorem 1, we have

$$\dim V_\chi^m(G) = \frac{n^{m-m}}{m} \sum_{d|m} S(h; m; d)n^d = \frac{1}{m} \sum_{d|m} \mu(m/d)n^d. \quad \square$$

Example 3 Let $G = \langle (12)(34)(5678) \rangle$ be a subgroup of \mathbb{S}_9 . Suppose χ is the identity character 1, i.e., $\chi = \chi_0$. Then by Theorem 1 we have

$$\begin{aligned} \dim V_\chi^9(G) &= \frac{n^{9-(2+2+4)}}{4} \sum_{\substack{d_1|2 \\ d_2|2 \\ d_3|4}} S(0; 2, 2, 4; d_1, d_2, d_3)n^{d_1+d_2+d_3} \\ &= \frac{n}{4} [S(0; 2, 2, 4; 2, 2, 4)n^8 + S(0; 2, 2, 4; 2, 2, 2)n^6 + S(0; 2, 2, 4; 1, 1, 1)n^3] \\ &= \frac{n}{4}[n^8 + n^6 + 2n^3]. \end{aligned}$$

References

- [1] T. M. Apostol, “*Introduction to Analytic Number Theory*”, Springer-Verlag, New York, 1976.
- [2] L. J. Cummings, *Cyclic Symmetry Classes*, J. Algebra **40** (1976), 401-405.
- [3] R. R. Holmes, T. Y. Tam, *Symmetry Classes of Tensors Associated with Certain Groups*, Linear and Multilinear Algebra **32** (1992), 21-31.
- [4] M. Marcus, “*Finite Dimensional Multilinear Algebra*”, Part 1, Marcel Dekker, New York, 1973.
- [5] R. Merris, *The Dimension of Certain Symmetry Classes of Tensors II*, Linear and Multilinear Algebra **4** (1976), 205-207.
- [6] R. Merris, M. A. Rashid, *The Dimensions of Certain Symmetry Classes of Tensors*, Linear and Multilinear Algebra **2** (1974), 245-248.
- [7] T. Y. Tam, *On the Cyclic Symmetry Classes*, J. Algebra **182** (1996), 557-560.

The Authors' Addresses

M. R. Darafsheh, Department of Mathematics and Computer Science, University of Tehran, and Institute for Studies in Theoretical Physics and Mathematics, Tehran, Iran.

E-mail address: darafshe@vax.ipm.ac.ir

M. R. Pournaki, Department of Mathematics and Computer Science, University of Tehran, and Institute for Studies in Theoretical Physics and Mathematics, Tehran, Iran.

E-mail address: pournaki@vax.ipm.ac.ir