

ERRATUM TO “ON THE EXISTENCE OF PERIODIC SOLUTIONS  
FOR A CLASS OF GENERALIZED FORCED LIÉNARD  
EQUATIONS” [APPL. MATH. LETT. 20 (2007), NO. 3, 248–254]

M. R. POURNAKI AND A. RAZANI

The authors would like to thank Professor Alessandro Fonda for pointing out the error in the statement of the Main Theorem. On pages 1 and 2 “ $T$ -periodic real function” should read “real function”. Also the statement of the Main Theorem should read as follows:

**Main Theorem.** *Suppose  $f$ ,  $k$ , and  $g$  are real functions on  $\mathbb{R}$  which are locally Lipschitz and  $p$  is a nonconstant, continuous, real function on  $[0, T]$ ,  $T > 0$ . Also suppose all solutions of the initial value problem (1.1) can be extended to  $[0, T]$ . If there exist real numbers  $a_1$  and  $a_2$  for which  $g(a_1) \leq p(t) \leq g(a_2)$  holds for each  $0 \leq t \leq T$ , then there exists  $T_0$ ,  $0 < T_0 < T$ , such that if  $p$  is  $T_0$ -periodic, Eq. (1.1) has at least one  $T_0$ -periodic solution.*

M. R. POURNAKI, DEPARTMENT OF MATHEMATICAL SCIENCES, SHARIF UNIVERSITY OF TECHNOLOGY, P.O. BOX 11155-9415, TEHRAN, IRAN, AND SCHOOL OF MATHEMATICS, INSTITUTE FOR STUDIES IN THEORETICAL PHYSICS AND MATHEMATICS, P.O. BOX 19395-5746, TEHRAN, IRAN.

*E-mail address:* [pournaki@ipm.ir](mailto:pournaki@ipm.ir)

*URL:* <http://math.ipm.ac.ir/pournaki/>

A. RAZANI, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, IMAM KHOMEINI INTERNATIONAL UNIVERSITY, P.O. BOX 34194-288, QAZVIN, IRAN, AND SCHOOL OF MATHEMATICS, INSTITUTE FOR STUDIES IN THEORETICAL PHYSICS AND MATHEMATICS, P.O. BOX 19395-5746, TEHRAN, IRAN.

*E-mail address:* [razani@ikiu.ac.ir](mailto:razani@ikiu.ac.ir)

*URL:* <http://math.ipm.ac.ir/razani/>