



On the h -vector of a simplicial complex with Serre's condition[☆]

A. Goodarzi^a, M.R. Pournaki^{b,c,*}, S.A. Seyed Fakhari^b, S. Yassemi^{a,c}

^a School of Mathematics, Statistics and Computer Science, College of Science, University of Tehran, Tehran, Iran

^b Department of Mathematical Sciences, Sharif University of Technology, P.O. Box 11155-9415, Tehran, Iran

^c School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box 19395-5746, Tehran, Iran

ARTICLE INFO

Article history:

Received 6 December 2010

Received in revised form 7 April 2011

Available online 25 June 2011

Communicated by A.V. Geramita

MSC:

Primary: 13F55; 05E45

Secondary: 13C14

ABSTRACT

Let Δ be a $(d - 1)$ -dimensional simplicial complex and let $h(\Delta) = (h_0, h_1, \dots, h_d)$ be its h -vector. A recent result of Murai and Terai guarantees that if Δ satisfies Serre's condition (S_r) , then (h_0, h_1, \dots, h_r) is an M -vector and $h_r + h_{r+1} + \dots + h_d$ is nonnegative. In this article, we extend the result of Murai and Terai by giving r extra necessary conditions. More precisely, we prove that if Δ satisfies Serre's condition (S_r) , then $\binom{i}{i} h_r + \binom{i+1}{i} h_{r+1} + \dots + \binom{i+d-r}{i} h_d$, $0 \leq i \leq r \leq d$, are all nonnegative.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction and preliminaries

Algebraic combinatorics is an area of mathematics that employs methods of abstract algebra in various combinatorial contexts. One of the fastest developing subfields within algebraic combinatorics is combinatorial commutative algebra. It has evolved into one of the most active areas of mathematics during the past several decades. We refer the reader to the book by Stanley [13] as a general reference in the subject.

The study of h -vectors of simplicial complexes has long been a topic of interest both in combinatorics and combinatorial commutative algebra (see, for example, [1, 12, 15, 3, 2, 17, 9, 14, 10, 11]). Let Δ be a $(d - 1)$ -dimensional simplicial complex and let $h(\Delta) = (h_0, h_1, \dots, h_d)$ be its h -vector. One of the fundamental questions in the subject is when h_0, \dots, h_d are all nonnegative. A classical result of Stanley guarantees that if Δ is Cohen–Macaulay, then h_i is nonnegative for every i with $0 \leq i \leq d$ (see [13, Theorem 3.3, Page 59]).

Recently, Murai and Terai [10] have generalized the classical result of Stanley in terms of Serre's condition, which appear in commutative algebra. They have proved that if Δ satisfies Serre's condition (S_r) , then (h_0, h_1, \dots, h_r) is an M -vector and $h_r + h_{r+1} + \dots + h_d$ is nonnegative. It is natural to ask whether these conditions are sufficient for a simplicial complex to satisfy Serre's condition. In fact, this is not the case, even for the independence simplicial complex of cycle graphs. In Theorem 2.1, we extend the result of Murai and Terai by giving r extra necessary conditions. These conditions are in fact sufficient for the independence simplicial complex of cycle graphs and quasi-forest simplicial complexes to satisfy Serre's condition (see Remark 2.2 and Corollary 2.5).

For convenience of the reader, let us recall some notions. Let $S = \mathbb{K}[x_1, \dots, x_n]$ be the polynomial ring in n variables over a field \mathbb{K} and let M be a nonzero finitely generated S -module. We say that M satisfies Serre's condition (S_r) , or simply M is a (S_r) module, if for every $\mathfrak{p} \in \text{Spec}(S)$, the inequality $\text{depth } M_{\mathfrak{p}} \geq \min\{r, \dim M_{\mathfrak{p}}\}$ holds true. It is easy to see that M is

[☆] The research of M.R. Pournaki and S. Yassemi was in part supported by a grant from IPM (No. 89130119 and No. 89130213).

* Corresponding author at: Department of Mathematical Sciences, Sharif University of Technology, P.O. Box 11155-9415, Tehran, Iran.

E-mail addresses: af.goodarzi@gmail.com (A. Goodarzi), pournaki@ipm.ir (M.R. Pournaki), fakhari@ipm.ir (S.A. Seyed Fakhari), yassemi@ipm.ir (S. Yassemi).

URLs: <http://math.ipm.ac.ir/pournaki/> (M.R. Pournaki), <http://math.ipm.ac.ir/fakhari/> (S.A. Seyed Fakhari), <http://math.ipm.ac.ir/yassemi/> (S. Yassemi).

Cohen–Macaulay if and only if it is a (S_r) module for all $r \geq 1$. We say that a simplicial complex Δ satisfies *Serre’s condition* (S_r) , or simply Δ is a (S_r) simplicial complex, if its Stanley–Reisner ring satisfies Serre’s condition (S_r) . Since every simplicial complex satisfies Serre’s condition (S_1) , we assume that $r \geq 2$. It is well known that if Δ is a (S_r) simplicial complex, then Δ is pure (see [10, Lemma 2.6]).

Let Δ be a simplicial complex. A facet F of Δ is called a *leaf* if there is a facet $G \neq F$ of Δ , called a *branch* of F , in such a way that $H \cap F \subseteq G \cap F$ holds true for all facets H of Δ with $H \neq F$. A *quasi-forest simplicial complex* is a simplicial complex Δ , which enjoys an ordering F_1, F_2, \dots, F_s of the facets of Δ , called a *leaf order*, in such a way that for every j with $1 \leq j \leq s$ the facet F_j is a leaf of the subcomplex $\langle F_1, \dots, F_j \rangle$. One can easily check that if Δ is a quasi-forest simplicial complex and $f(\Delta) = (f_0, f_1, \dots, f_{d-1})$ is its f -vector, then $f_{d-1} \leq n - d + 1$ and the equality holds if and only if Δ is Cohen–Macaulay.

2. The main result

In this section, we state and prove the main result of this article. Recently, Murai and Terai [10] have proved that if Δ is a $(d - 1)$ -dimensional (S_r) simplicial complex and $h(\Delta) = (h_0, h_1, \dots, h_d)$ is its h -vector, then the following two conditions hold:

- (h_0, h_1, \dots, h_r) is an M -vector, and
- $h_r + h_{r+1} + \dots + h_d$ is nonnegative.

Note that in the above, a sequence of integers $\mathfrak{h} = (h_0, h_1, \dots, h_r)$ is called an M -vector provided there exists a Cohen–Macaulay $(r - 1)$ -dimensional simplicial complex Γ with $h(\Gamma) = \mathfrak{h}$. Now, it is natural to ask whether the above two conditions are sufficient for a simplicial complex to be (S_r) . This is not the case, even for the independence simplicial complex of cycle graphs, as C_9 shows. We now extend the result of Murai and Terai by giving r extra necessary conditions to obtain sufficient conditions for the independence simplicial complex of cycle graphs as well as for quasi-forest simplicial complexes to be (S_r) (see Remark 2.2 and Corollary 2.5). Note that condition (2) in Theorem 2.1 includes $r + 1$ statements and it gives us the second condition of Murai and Terai for $i = 0$.

Theorem 2.1. *Let Δ be a $(d - 1)$ -dimensional (S_r) simplicial complex and let $h(\Delta) = (h_0, h_1, \dots, h_d)$ be its h -vector. Then the following conditions hold:*

- (1) (h_0, h_1, \dots, h_r) is an M -vector, and
- (2) $\binom{i}{j} h_r + \binom{i+1}{j} h_{r+1} + \dots + \binom{i+d-r}{j} h_d$ is nonnegative for every i with $0 \leq i \leq r \leq d$.

Proof. Condition (1) was proved in [10]. We prove condition (2) by using induction on i . If $i = 0$, then condition (2) reduces to

$$h_r + h_{r+1} + \dots + h_d \geq 0,$$

which is true by Theorem 1.3 of [10]. Let $1 \leq i \leq r$ be an arbitrary integer. We now suppose that condition (2) is true for $i - 1$ and prove that it holds true for i .

In order to do this, note that if $d < r$, then there is nothing to prove. Therefore, we assume that $d \geq r$ and we use induction on d . If $d = r$, then we must show that h_r is nonnegative and this is true since Δ is Cohen–Macaulay (see [13, Theorem 3.3, Page 59]). We now suppose that $d > r$. By [15, Proposition 2.3], we have the following identity, where $h_i(\text{lk}_\Delta(v))$, $0 \leq i \leq d - 1$, denotes the i th component of the h -vector of the simplicial complex $\text{lk}_\Delta(v)$ and Δ is assumed on $[n] = \{1, \dots, n\}$:

$$ih_i + (d - i + 1)h_{i-1} = \sum_{v \in [n]} h_{i-1}(\text{lk}_\Delta(v)).$$

Therefore, we have

$$\begin{aligned} (r + 1)h_{r+1} + (d - r)h_r &= \sum_{v \in [n]} h_r(\text{lk}_\Delta(v)), \\ (r + 2)h_{r+2} + (d - r - 1)h_{r+1} &= \sum_{v \in [n]} h_{r+1}(\text{lk}_\Delta(v)), \\ &\vdots \\ dh_d + h_{d-1} &= \sum_{v \in [n]} h_{d-1}(\text{lk}_\Delta(v)). \end{aligned}$$

Now, by multiplying the j th equation by $\binom{i+j-1}{i}$, $1 \leq j \leq d - r - 1$, and then adding the resulting equations, we conclude that

$$(d - r)h_r + ((i + 1)d - i(r + 1))h_{r+1} + \dots + d \binom{i + d - r - 1}{i} h_d = \sum_{j=0}^{d-r-1} \binom{i+j}{i} \sum_{v \in [n]} h_{r+j}(\text{lk}_\Delta(v)).$$

Since, by Lemma 2.2 of [4], $\text{lk}_\Delta(v)$ is a (S_r) simplicial complex for every vertex v of Δ , by the induction hypothesis on d ,

$$\sum_{j=0}^{d-r-1} \binom{i+j}{i} \sum_{v \in [n]} h_{r+j}(\text{lk}_\Delta(v)) \geq 0,$$

and therefore we have

$$(d-r)h_r + ((i+1)d - i(r+1))h_{r+1} + \dots + d \binom{i+d-r-1}{i} h_d \geq 0.$$

On the other hand, by the induction hypothesis on i , we have

$$\binom{i-1}{i-1} h_r + \binom{i}{i-1} h_{r+1} + \dots + \binom{i+d-r-1}{i-1} h_d \geq 0.$$

Multiplying the last inequality by $r-i$ and adding the result to the previous inequality and then dividing by $d-i$ implies that

$$\binom{i}{i} h_r + \binom{i+1}{i} h_{r+1} + \dots + \binom{i+d-r}{i} h_d \geq 0.$$

Therefore, condition (2) holds true by induction. \square

Let C_n be the cycle graph with n vertices ($n \geq 3$). It is known that the independence simplicial complex of C_n is Cohen–Macaulay if and only if $n = 3, 5$ (see [16, Corollary 4.7]). Also, the independence simplicial complex of C_n is (S_2) if and only if $n = 3, 5, 7$ (see [5, Proposition 1.6]). Similarly, the independence simplicial complex of C_n is (S_r) for $r \geq 3$ if and only if $n = 3, 5$ (see [4, Proposition 4.2]). These observations together with an easy computation imply the following remark.

Remark 2.2. Let (h_0, h_1, \dots, h_d) be the h -vector of the independence simplicial complex of C_n . Then C_n is Cohen–Macaulay if and only if h_0, \dots, h_d are all nonnegative. Also C_n is (S_r) for $r \geq 2$ if and only if the following conditions hold:

- (1) h_0, \dots, h_{r-1} are all nonnegative,
- (2) $h_r + h_{r+1} + \dots + h_d$ is nonnegative, and
- (3) $h_r + 2h_{r+1} + \dots + (d-r+1)h_d$ is nonnegative.

Therefore, conditions (1) and (2) of Theorem 2.1 are equivalent to C_n being (S_r) .

Recently, the concept of quasi-forest simplicial complex has been extensively studied in combinatorics and combinatorial commutative algebra (see, for example, [8,18,6]). In the following we show that nonnegativity of the components of the h -vector of a quasi-forest simplicial complex is sufficient for Cohen–Macaulayness of it.

Proposition 2.3. Let Δ be a $(d-1)$ -dimensional quasi-forest simplicial complex and let $h(\Delta) = (h_0, h_1, \dots, h_d)$ be its h -vector. Then Δ is Cohen–Macaulay if and only if for every i with $0 \leq i \leq d$ we have $h_i \geq 0$.

Proof. If Δ is Cohen–Macaulay, then by [13, Theorem 3.3, Page 59], for every i with $0 \leq i \leq d$, we have $h_i \geq 0$.

Conversely, suppose that for every i with $0 \leq i \leq d$, we have $h_i \geq 0$. Let $f(\Delta) = (f_0, f_1, \dots, f_{d-1})$ be the f -vector of Δ and $h(\Delta) = (h_0, h_1, \dots, h_d)$ be the h -vector of Δ . Since Δ is a quasi-forest simplicial complex, $f_{d-1} \leq n-d+1$. Therefore, we conclude that $h_2 + h_3 + \dots + h_d \leq 0$ and so by assumption, for every i with $2 \leq i \leq d$, we have $h_i = 0$. Now Theorem B of [7] implies that Δ is Cohen–Macaulay. \square

In [5], it is shown that for the independence simplicial complex of bipartite and chordal graphs, Cohen–Macaulayness is equivalent to the Serre’s condition (S_2) . In the following proposition, we show that the same result holds true for quasi-forest simplicial complexes.

Proposition 2.4. Let Δ be a $(d-1)$ -dimensional quasi-forest simplicial complex. Then Δ is Cohen–Macaulay if and only if Δ satisfies Serre’s condition (S_2) .

Proof. If Δ is Cohen–Macaulay, then obviously Δ satisfies Serre’s condition (S_2) .

Conversely, suppose that Δ satisfies Serre’s condition (S_2) . Let $f(\Delta) = (f_0, f_1, \dots, f_{d-1})$ be the f -vector of Δ and $h(\Delta) = (h_0, h_1, \dots, h_d)$ be the h -vector of Δ . Suppose, on the contrary, that Δ is not Cohen–Macaulay. We have $f_{d-1} \leq n-d$ and so $h_0 + h_1 + (h_2 + \dots + h_d) \leq n-d$. Thus $h_2 + \dots + h_d$ is negative, a contradiction. Therefore Δ is Cohen–Macaulay. \square

Corollary 2.5. Let Δ be a $(d-1)$ -dimensional quasi-forest simplicial complex and let $h(\Delta) = (h_0, h_1, \dots, h_d)$ be its h -vector. Then Δ satisfies Serre’s condition (S_2) if and only if $h_2 + h_3 + \dots + h_d$ is nonnegative.

Proof. If Δ satisfies Serre’s condition (S_2) , then by Theorem 1.1 of [10], $h_2 + h_3 + \dots + h_d$ is nonnegative.

Conversely, assume that $h_2 + h_3 + \dots + h_d$ is nonnegative. Let $f(\Delta) = (f_0, f_1, \dots, f_{d-1})$ be the f -vector of Δ and $h(\Delta) = (h_0, h_1, \dots, h_d)$ be the h -vector of Δ . Suppose, on the contrary, that Δ does not satisfies Serre’s condition (S_2) . Then by Proposition 2.4, Δ is not Cohen–Macaulay and so $f_{d-1} \leq n-d$. Thus $h_2 + h_3 + \dots + h_d < 0$, a contradiction. Therefore Δ satisfies Serre’s condition (S_2) . \square

Although, as we mentioned earlier, the conditions of Murai and Terai in [10] are not sufficient for a simplicial complex to be (S_r) , even for the independence simplicial complex of a cycle graph, our conditions (1) and (2) stated in [Theorem 2.1](#) are sufficient for the independence simplicial complex of cycle graphs as well as for quasi-forest simplicial complexes to be (S_r) (see [Remark 2.2](#) and [Corollary 2.5](#)). Now, it is natural to ask whether the converse of [Theorem 2.1](#) is true in general. We close the article by stating the following question which is a converse to [Theorem 2.1](#) in a standard sense.

Question 2.6. *Let d and r be integers with $d \geq r \geq 2$ and let $\mathfrak{h} = (h_0, h_1, \dots, h_d)$ be the h -vector of a simplicial complex in such a way that the following conditions hold:*

- (1) (h_0, h_1, \dots, h_r) is an M -vector, and
- (2) $\binom{d}{i}h_r + \binom{d+1}{i}h_{r+1} + \dots + \binom{d+r-i}{i}h_d$ is nonnegative for every i with $0 \leq i \leq r \leq d$.

Does there exist a $(d - 1)$ -dimensional (S_r) simplicial complex Δ with $h(\Delta) = \mathfrak{h}$?

Acknowledgements

The authors would like to thank Satoshi Murai and Naoki Terai for reading an earlier version of this article and for their useful comments.

References

- [1] A. Björner, P. Frankl, R.P. Stanley, The number of faces of balanced Cohen–Macaulay complexes and a generalized Macaulay theorem, *Combinatorica* 7 (1) (1987) 23–34.
- [2] W. Bruns, T. Römer, h -Vectors of Gorenstein polytopes, *J. Combin. Theory Ser. A* 14 (1) (2007) 65–76.
- [3] V.E. de Quehen, L.G. Roberts, Non-Cohen–Macaulay projective monomial curves with positive h -vector, *Canad. Math. Bull.* 48 (2) (2005) 203–210.
- [4] H. Haghighi, N. Terai, S. Yassemi, R. Zaare-Nahandi, Sequentially S_r simplicial complexes and sequentially S_2 graphs, *Proc. Amer. Math. Soc.* 139 (6) (2011) 1993–2005.
- [5] H. Haghighi, S. Yassemi, R. Zaare-Nahandi, Bipartite S_2 graphs are Cohen–Macaulay, *Bull. Math. Soc. Sci. Math. Roumanie (NS)* 53(101) (2) (2010) 125–132.
- [6] J. Herzog, T. Hibi, S. Murai, N.V. Trung, X. Zheng, Kruskal–Katona type theorems for clique complexes arising from chordal and strongly chordal graphs, *Combinatorica* 28 (3) (2008) 315–323.
- [7] J. Herzog, T. Hibi, S. Murai, N.V. Trung, X. Zheng, Kruskal–Katona type theorems for clique complexes arising from chordal and strongly chordal graphs, [arXiv:math/0606477v1](https://arxiv.org/abs/math/0606477v1).
- [8] J. Herzog, T. Hibi, X. Zheng, Dirac’s theorem on chordal graphs and Alexander duality, *European J. Combin.* 25 (7) (2004) 949–960.
- [9] J. Migliore, U. Nagel, F. Zanello, On the degree two entry of a Gorenstein h -vector and a conjecture of Stanley, *Proc. Amer. Math. Soc.* 136 (8) (2008) 2755–2762.
- [10] S. Murai, N. Terai, h -Vectors of simplicial complexes with Serre’s conditions, *Math. Res. Lett.* 16 (6) (2009) 1015–1028.
- [11] J. Schweig, On the h -vector of a lattice path matroid, *Electron. J. Combin.* 17 (1) (2010) Note 3, 6 pp.
- [12] R.P. Stanley, f -Vectors and h -vectors of simplicial posets, *J. Pure Appl. Algebra* 71 (2–3) (1991) 319–331.
- [13] R.P. Stanley, *Combinatorics and Commutative Algebra*, Second edition, in: *Progress in Mathematics*, 41, Birkhäuser Boston, Inc., Boston, MA, 1996.
- [14] J.R. Stembridge, Coxeter cones and their h -vectors, *Adv. Math.* 217 (5) (2008) 1935–1961.
- [15] E. Swartz, Lower bounds for h -vectors of k -CM, independence, and broken circuit complexes, *SIAM J. Discrete Math.* 18 (3) (2004/05) 647–661.
- [16] R.H. Villarreal, Cohen–Macaulay graphs, *Manuscripta Math.* 66 (3) (1990) 277–293.
- [17] F. Zanello, The h -vector of a relatively compressed level algebra, *Comm. Algebra* 35 (4) (2007) 1087–1091.
- [18] X. Zheng, Resolutions of facet ideals, *Comm. Algebra* 32 (6) (2004) 2301–2324.