

NEW CLASSES OF SET-THEORETIC COMPLETE INTERSECTION MONOMIAL IDEALS

M. R. Pournaki^{1,2}, S. A. Seyed Fakhari², and S. Yassemi^{2,3}

¹Department of Mathematical Sciences, Sharif University of Technology, Tehran, Iran

²School of Mathematics, Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

³School of Mathematics, Statistics and Computer Science, College of Science, University of Tehran, Tehran, Iran

Let Δ be a simplicial complex and χ be an s -coloring of Δ . Biermann and Van Tuyl have introduced the simplicial complex Δ_χ . As a corollary of Theorems 5 and 7 in their 2013 article, we obtain that the Stanley–Reisner ring of Δ_χ over a field is Cohen–Macaulay. In this note, we generalize this corollary by proving that the Stanley–Reisner ideal of Δ_χ over a field is set-theoretic complete intersection. This also generalizes a result of Macchia.

Key Words: Arithmetical rank; Lyubeznik resolution; s -Coloring; Set-theoretic complete intersection ideal; Simplicial complex.

2010 Mathematics Subject Classification: 13F55; 13A15.

1. STATEMENT OF THE MAIN THEOREM

Let us start this note with some notions of combinatorics. A *simplicial complex* Δ on the set of *vertices* $V = \{v_1, \dots, v_n\}$ is a collection of subsets of V which is closed under taking subsets; that is, if $F \in \Delta$ and $F' \subseteq F$, then also $F' \in \Delta$. Every element $F \in \Delta$ is called a *face* of Δ , and a *facet* of Δ is a maximal face of Δ with respect to inclusion. It is clear that all facets of Δ determines it. When F_1, \dots, F_t are all facets of Δ , we write $\Delta = \langle F_1, \dots, F_t \rangle$. In this case, we say χ is an s -coloring of Δ when χ is a partition of V , say $V = W_1 \cup \dots \cup W_s$ (where the sets W_j are allowed to be empty), such that for every $1 \leq i \leq t$ and every $1 \leq j \leq s$ the inequality $|F_i \cap W_j| \leq 1$ holds true. Biermann and Van Tuyl [3] have defined a new simplicial complex Δ_χ on the set of vertices $\{v_1, \dots, v_n, w_1, \dots, w_s\}$ with faces $\sigma \cup \tau$, where σ is a face of Δ and τ is any subset of $\{w_1, \dots, w_s\}$ such that for every $w_j \in \tau$, we have $\sigma \cap W_j = \emptyset$. It is shown in [3] that the facets of Δ_χ are in the form $F \cup F'$, where F is any face of Δ and $F' = \{w_i \mid F \cap W_i = \emptyset\}$. The simplicial complex Δ_χ is generally larger than Δ . For example, if $\Delta = \langle \{v_1, v_2, v_3\}, \{v_3, v_4\}, \{v_4, v_5, v_6\} \rangle$ is the simplicial complex on

Received March 28, 2014; Revised May 28, 2014. Communicated by G. Leuschke.

Address correspondence to Dr. M. R. Pournaki, Department of Mathematical Sciences, Sharif University of Technology, P.O. Box 11155-9415, Tehran, Iran; E-mail: pournaki@ipm.ir

the set of vertices $V = \{v_1, v_2, v_3, v_4, v_5, v_6\}$ and χ is the 3-coloring of Δ given by $V = W_1 \cup W_2 \cup W_3$, where $W_1 = \{v_1, v_4\}$, $W_2 = \{v_2, v_5\}$, and $W_3 = \{v_3, v_6\}$, then we have

$$\Delta_\chi = \left\langle \{w_1, w_2, w_3\}, \{v_1, w_2, w_3\}, \{v_2, w_1, w_3\}, \{v_3, w_1, w_2\}, \right. \\ \{v_4, w_2, w_3\}, \{v_5, w_1, w_3\}, \{v_6, w_1, w_2\}, \{v_1, v_2, w_3\}, \\ \{v_1, v_3, w_2\}, \{v_2, v_3, w_1\}, \{v_3, v_4, w_2\}, \{v_4, v_5, w_3\}, \\ \left. \{v_4, v_6, w_2\}, \{v_5, v_6, w_1\}, \{v_1, v_2, v_3\}, \{v_4, v_5, v_6\} \right\rangle.$$

Although many properties of the simplicial complex Δ_χ are known, it might be complicated in some sense comparing with Δ .

We now switch to the connection of the above-mentioned notions with commutative algebra. Let Δ be a simplicial complex on the set of vertices $V = \{v_1, \dots, v_n\}$. The Stanley–Reisner ideal of Δ over a field \mathbb{K} is the ideal I_Δ of $R = \mathbb{K}[x_1, \dots, x_n]$, the polynomial ring in n variables over \mathbb{K} , which is defined as follows:

$$I_\Delta = \left\langle \prod_{v_i \in F} x_i \mid F \notin \Delta \right\rangle.$$

Also, the Stanley–Reisner ring of Δ over \mathbb{K} is defined as R/I_Δ . One may ask about the arithmetical rank of the Stanley–Reisner ideal of a simplicial complex and whether it being set-theoretic complete intersection. We recall that for a given unitary commutative Noetherian ring R and for an ideal I of R , the arithmetical rank of I is denoted by $\text{ara}(I)$ and is defined as the smallest integer t for which there exist $a_1, \dots, a_t \in R$ such that $\sqrt{\langle a_1, \dots, a_t \rangle} = \sqrt{I}$. The inequality $\text{ht}(I) \leq \text{ara}(I)$ holds true generally and in the case of equality, I is called set-theoretic complete intersection. The arithmetical rank of monomial ideals of $R = \mathbb{K}[x_1, \dots, x_n]$ has been studied by several authors (see, for example, [1, 2, 6, 8, 9]).

We are now in the position to state the main theorem of this note. In the next section, we give a proof of the main theorem.

Main Theorem. *Let Δ be a simplicial complex and χ be an s -coloring of Δ . Then the Stanley–Reisner ideal of Δ_χ over a field is set-theoretic complete intersection.*

Before closing this section, we would like to point out that our main theorem generalizes two results of [3] and [9]. Let us explain some more things in detail to see this.

The Main Theorem as a Generalization of a Result of [3] Let $R = \mathbb{K}[x_1, \dots, x_n]$ and I be a squarefree monomial ideal of R . Lyubeznik [8] has proven that $\text{pd}_R(R/I) \leq \text{ara}(I)$, where $\text{pd}_R(R/I)$ denotes the projective dimension of R/I . Thus

$$\text{ht}(I) \leq \text{pd}_R(R/I) \leq \text{ara}(I).$$

This implies that, if I is set-theoretic complete intersection, then R/I is Cohen–Macaulay. Combining the above observation with our main theorem gives the following corollary, which is also a corollary of Theorems 5 and 7 in the article [3] of Biermann and Van Tuyl.

Corollary I ([3, Theorems 5 and 7]). *Let Δ be a simplicial complex and χ be an s -coloring of Δ . Then the Stanley–Reisner ring of Δ_χ over a field is Cohen–Macaulay.*

The Main Theorem as a Generalization of a Result of [9] Let G be a finite simple graph with the set of vertices $V(G) = \{v_1, \dots, v_n\}$. To the graph G , one can associate an ideal $I(G) \subseteq \mathbb{K}[x_1, \dots, x_n]$ generated by all monomials $x_i x_j$ such that v_i and v_j are adjacent. The ideal $I(G)$ is called the *edge ideal* of G over \mathbb{K} . We recall that $A \subseteq V(G)$ is an *independent set* in G if none of its elements are adjacent. Based on this notion, the *independence simplicial complex* of G is defined by

$$\Delta_G = \{A \subseteq V(G) \mid A \text{ is an independent set in } G\}.$$

Note that Δ_G is precisely the simplicial complex with the Stanley–Reisner ideal $I(G)$. Cook and Nagel [4] have defined the fully clique-whiskered graphs in the following way. For a given graph G , $C \subseteq V(G)$ is called a *clique* of G if every pair of vertices of C are adjacent in G . Let χ be a partition of $V(G)$, say $V(G) = W_1 \cup \dots \cup W_s$, such that W_i is a clique of G for every $1 \leq i \leq s$. Add new vertices y_1, \dots, y_s and new edges vy_i for every $v \in W_i$ and every $1 \leq i \leq s$. The resulting graph is called a *fully clique-whiskered graph* of G . One can easily see that the independence simplicial complex of this latter graph is Δ_χ and so our main theorem gives the following corollary, which is Proposition 4.1 in the article [9] of Macchia.

Corollary II ([9, Proposition 4.1]). *Let G be a finite simple graph. Then the edge ideals of fully clique-whiskered graphs of G over a field are all set-theoretic complete intersections.*

2. PROOF OF THE MAIN THEOREM

We are now ready to prove the main theorem. In order to do this, suppose that the set of vertices of Δ is $V = \{v_1, \dots, v_n\}$ and $V = W_1 \cup \dots \cup W_s$ is the s -coloring of Δ given by χ . Also, let $R = \mathbb{K}[x_1, \dots, x_n]$ be the polynomial ring in n variables over a field \mathbb{K} . Suppose that $\{v_1, \dots, v_n, w_1, \dots, w_s\}$ is the set of vertices of Δ_χ , and consider the Stanley–Reisner ideal I_{Δ_χ} of Δ_χ over \mathbb{K} as a monomial ideal in the polynomial ring $R' = \mathbb{K}[x_1, \dots, x_n, y_1, \dots, y_s]$.

Lemma A. *We have $\text{ht}(I_{\Delta_\chi}) = n$.*

Proof. One can easily see that

$$I_{\Delta_\chi} = I_\Delta + \langle x_i y_j \mid v_i \in W_j, 1 \leq j \leq s \rangle.$$

Note that the ideal $\langle x_1, \dots, x_n \rangle$ is a minimal prime ideal of I_{Δ_χ} . Now [3, Theorem 5] implies that $\text{ht}(I_{\Delta_\chi}) = n$, as required. \square

Let us introduce the Lyubeznik resolution, which is defined in [7]. For every monomial ideal I of a polynomial ring R , there is an explicit free resolution of R/I , which is defined as follows. Assume that $G(I) = \{m_1, \dots, m_p\}$ is the minimal set of monomial generators of I . For every integer $t \geq 1$ and every sequence $1 \leq \ell_1 < \dots < \ell_t \leq p$, consider the symbol $e_{\ell_1 \dots \ell_t}$. Let T_t be the free R -module generated by all symbols $e_{\ell_1 \dots \ell_t}$, $1 \leq \ell_1 < \dots < \ell_t \leq p$. The differential map d_t is also defined by

$$d_t(e_{\ell_1 \dots \ell_t}) = \sum_{k=1}^t (-1)^{k-1} \frac{\text{lcm}(m_{\ell_1}, \dots, m_{\ell_t})}{\text{lcm}(m_{\ell_1}, \dots, \widehat{m_{\ell_k}}, \dots, m_{\ell_t})} e_{\ell_1 \dots \widehat{\ell_k} \dots \ell_t}.$$

Then the complex (T_\bullet, d_\bullet) is a free resolution of R/I and is called the *Taylor resolution* of I (see, for example, [5, Theorem 7.1.1]). Now fix an order $m_1 > \dots > m_p$ on the set of minimal monomial generators of I . The symbol $e_{\ell_1 \dots \ell_t}$ is called *L-admissible* if m_q does not divide $\text{lcm}(m_{\ell_k}, \dots, m_{\ell_t})$ for every $1 \leq k < t$ and every $1 \leq q < \ell_k$. Then the *Lyubeznik resolution* of I with respect to the above order of the minimal monomial generators is a subcomplex of the Taylor resolution generated by all *L-admissible* symbols. Since being *L-admissible* depends on an order of minimal monomial generators of I , a Lyubeznik resolution also depends on it. Finally, we say that an *L-admissible* symbol $e_{\ell_1 \dots \ell_t}$ is *maximal* if $e_{k_1 \dots k_r}$ is not *L-admissible* whenever $\{\ell_1, \dots, \ell_t\} \not\supseteq \{k_1, \dots, k_r\}$.

Lemma B. *There is a Lyubeznik resolution of I_{Δ_χ} with length $\leq n$.*

Proof. Let $>$ be the pure lexicographic order induced by $x_1 > \dots > x_n > y_1 > \dots > y_s$ (see [5, Example 2.1.2(c)]), and consider the Lyubeznik resolution of I_{Δ_χ} induced by $>$. We show that this specific Lyubeznik resolution of I_{Δ_χ} has length $\leq n$. In order to do this, let $m_1 > \dots > m_p$ be the minimal monomial generators of I_{Δ_χ} and $e_{\ell_1 \dots \ell_k}$ be a maximal *L-admissible* symbol. We use induction on n to show that $k \leq n$. This completes the proof of the lemma.

One can easily check that $k \leq n$ is true for $n = 1, 2$. Thus we may assume that $n \geq 3$. Note that there exists a unique integer, say i_0 , with $1 \leq i_0 \leq s$, such that $v_1 \in W_{i_0}$. We claim that there exists an integer $1 \leq i \leq n$ such that x_i divides m_{ℓ_1} but does not divide $\text{lcm}(m_{\ell_2}, \dots, m_{\ell_k})$. In order to prove the claim, since $e_{\ell_1 \dots \ell_k}$ is a maximal *L-admissible* symbol, by using [6, Lemma 3], it follows that $\ell_1 = 1$. This implies that x_1 divides m_{ℓ_1} . Now, by the definition of *L-admissible* symbol, we conclude that there is a variable which divides m_{ℓ_1} but does not divide $\text{lcm}(m_{\ell_2}, \dots, m_{\ell_k})$. If all of the variables which divide m_{ℓ_1} belong to the set $\{x_1, \dots, x_n\}$, then we are done. Otherwise, we have $m_{\ell_1} = x_1 y_{i_0}$ since $x_1 \mid m_{\ell_1}$ and $v_1 \in W_{i_0}$. By the choice of $>$, we conclude that x_1 does not divide $\text{lcm}(m_{\ell_2}, \dots, m_{\ell_k})$, and so the claim holds true. By the claim, we may assume that x_i divides m_{ℓ_1} but does not divide $\text{lcm}(m_{\ell_2}, \dots, m_{\ell_k})$. Let

$$\Delta' = \Delta \setminus \{v_i\} = \{F \in \Delta \mid v_i \notin F\}$$

be a simplicial complex on the set of vertices $V' = V \setminus \{v_i\}$ and χ' be an s -coloring of Δ' given by $V' = W'_1 \cup \dots \cup W'_s$, where $W'_j = W_j \setminus \{v_i\}$ for every $1 \leq j \leq s$. It is clear that $m_{\ell_2}, \dots, m_{\ell_k}$ belong to the set of minimal monomial generators of $I_{\Delta'_\chi}$ and they

determine an L -admissible symbol of $I_{\Delta'_\lambda}$. Since Δ' has $n - 1$ vertices, our induction hypothesis implies that $k - 1 \leq n - 1$ and thus $k \leq n$. \square

We also need to use the following known result of Kimura.

Lemma C ([6, Theorem 1]). *Let I be a monomial ideal of a polynomial ring and λ be the minimum length of the Lyubeznik resolutions of I . Then $\text{ara}(I) \leq \lambda$.*

Now, by combining Lemmas B and C, we get $\text{ara}(I_{\Delta_\lambda}) \leq n$. Therefore, Lemma A implies that

$$n = \text{ht}(I_{\Delta_\lambda}) \leq \text{pd}_R(R/I_{\Delta_\lambda}) \leq \text{ara}(I_{\Delta_\lambda}) \leq n,$$

and so $\text{ht}(I_{\Delta_\lambda}) = \text{ara}(I_{\Delta_\lambda}) = n$. This means that I_{Δ_λ} is a set-theoretic complete intersection ideal of R' and completes the proof of the main theorem. \square

ACKNOWLEDGMENT

The authors would like to thank the referee for a careful reading of this note and for valuable comments.

FUNDING

The research of M. R. Pournaki, S. A. Seyed Fakhari, and S. Yassemi was in part supported by a grant from IPM (No. 93130115, No. 93130422, and No. 93130214).

REFERENCES

- [1] Barile, M. (2008). On the arithmetical rank of the edge ideals of forests. *Comm. Algebra* 36(12):4678–4703.
- [2] Barile, M., Kiani, D., Mohammadi, F., Yassemi, S. (2012). Arithmetical rank of the cyclic and bicyclic graphs. *J. Algebra Appl.* 11(2):1250039, 14 pp.
- [3] Biermann, J., Van Tuyl, A. (2013). Balanced vertex decomposable simplicial complexes and their h -vectors. *Electron. J. Combin.* 20(3): Paper 15, 12 pp.
- [4] Cook, D., Nagel, U. (2012). Cohen–Macaulay graphs and face vectors of flag complexes. *SIAM J. Discrete Math.* 26(1):89–101.
- [5] Herzog, J., Hibi, T. (2011). *Monomial Ideals*. London: Springer-Verlag.
- [6] Kimura, K. (2009). Lyubeznik resolutions and the arithmetical rank of monomial ideals. *Proc. Amer. Math. Soc.* 137(11):3627–3635.
- [7] Lyubeznik, G. (1988). A new explicit finite free resolution of ideals generated by monomials in an R -sequence. *J. Pure Appl. Algebra* 51(1–2):193–195.
- [8] Lyubeznik, G. (1988). On the arithmetical rank of monomial ideals. *J. Algebra* 112(1):86–89.
- [9] Macchia, A. (2015). The arithmetical rank of the edge ideals of graphs with whiskers. *Beitr. Algebra Geom.* 56(1):147–158.